

Bortrac 150

RECOMMENDED APPLICATION RATES AND TIMINGS

Almond, Nuts (Deciduous): 1 L/ha at spring bud burst, first emergent leaves and again during nut development. Alternatively, 1 L/ha at bud break and 2 L/ha after harvest before senescence. Water rate: 500 to 1000 L/ha.

Apple, Pears: 1-2 L/ha applied at pink bud, start of flowering, and again at petal fall. 2 L/ha after harvest but before leaf senescence. Water rate: 500-1000 L/ha.

Apricot, Nectarines, Peach, Plum: 1 to 2 L/ha at winter bud and again at pink bud. 2 L/ha after harvest but before leaf senescence. Water rate: 500-1000 L/ha.

Avocado: 2 applications of 1.5-2.5 L/ha applied at spring bud development and again at spring flush. Water rate: 500-1000 L/ha

Bananas: Regular applications may be necessary where low or marginal levels of boron exist. Spray at 300 mL-1 L/ha per application. Water rate: 30-500 L/ha.

Brassicas, Canola: 2-3 L/ha applied at 4 to 9 leaf stage then at onset of stem extension or prior to head development. Water rate: 50-500 L/ha.

Citrus: 2-3 L/ha at white bud or when flower buds separated. Water rate: 500-1000 L/ha.

Cucumbers, Cucurbits Pumpkin, Squash, Water Melon, Zucchini (Field Grown): 2 L/ha at 4 leaf stage, repeated if necessary 10 to 14 days later. Water rate: 50-200 L/ha.

Grapevines: Soil application: 3 to 5 L/ha. Foliar application: 1.5 L/ha at flower truss visible and at flower buds separated, and again at fruit set. Water rate: 500-1000 L/ha.

Mango: One or two applications of 2 L/ha at bud development to early flush at a 10 to 14 day interval. Water rate: 500-1000 L/ha.

Melon (field grown): 2 L/ha at 4 leaf stage, repeated if necessary 10 to 14 days later. Water rate: 50-200 L/ha.

Olive: 2 to 3 L/ha applied at bud break and again at the opening of the calyx if necessary. In addition, one or two applications of 2 to 3 L/ha at a 10 to 14 day interval post-flowering. Also, 2 to 3 L/ha applied post-harvest. Water rate: 400-1000 L/ha.

Potatoes: 2 L/ha 10 to 14 days after 100% emergence. Repeat if necessary 10 to 14 days later. Water rate: 50-200 L/ha.

Strawberry (field grown): 1.5 L/ha applied at green bud and again at white bud. Water rate: 200-500 L/ha.

Tomato (Field Grown): 2 L/ha when plants are 15 cm high. Repeat if necessary at 10 day intervals. Water rate: 50-500 L/ha.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS.

Mix the product thoroughly before use. Pour through the top filter into the tank three-quarters filled with water, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use and advice regarding whether or not product(s) should be co-applied. Many variables, outside the control of Yara, can influence the performance of co-applied products and therefore co-application is entirely at the risk of the end-user. It is strongly recommended that a limited application is made initially when using unfamiliar product mixes. *Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.*

PRECAUTIONS


Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. *Protect from frost and other climatic extremes. (temperatures exceeding 40°C)*

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY

Yara warrants that the product within the unopened container conforms to the chemical description on this label. The directions on this label must be followed carefully and are based on tests, believed to be reliable. As the occurrence of unusual weather or soil conditions, and the use of this product, are all beyond the control of Yara, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of Yara exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Goods are sold subject to the standard Conditions of Sale of Yara, available on request or at www.yaravita.com. Use of the product implies that the purchaser accepts these Conditions of Sale.



5

For further crop recommendations visit Product Information at www.yara.com.au