

Hydromag Liquid

RECOMMENDED APPLICATION RATES AND TIMINGS

Apple, Pears: 3-4 L/ha applied at pink bud and again at fruit set*.

In case of moderate or severe deficiency repeat during fruit development*. 5 L/ha after harvest but before leaf senescence.

Water rate: 500-1000 L/ha. *For application during fruit set and fruit development, it may, under certain conditions, be necessary to delay applications on varieties sensitive to russetting.

Apricot, Cherry, Nectarines, Peach, Plum: 4 L/ha during fruit development. 5 L/ha after harvest but before leaf senescence.

Water rate: 500-1000 L/ha.

Avocado: 4-6 L/ha applied during spring flush and again during summer flush. Water rate: 500-1000 L/ha.

Bananas: Regular applications may be necessary where low to marginal magnesium levels exist. Spray at 2-8 L/ha per application

Water rate: 30-500 L/ha.

Brassicac, Canola: 2-3 L/ha commencing at 4 to 9 true leaves. Water rate: 50-500 L/ha.

Cereals: 1.5-4 L/ha applied during tillering and again at flag leaf emergence. Water rate: 50-500 L/ha.

Citrus: Two applications of 3 L/ha applied during spring flush and again during autumn flush. Water rate: 500-1000 L/ha.

Cucumber, Cucurbits, Pumpkin, Squash, Water Melon, Zucchini (Field Grown): 2 to 3 applications of 4 L/ha commencing when sufficient leaf area to intercept spray. Water rate: 50-200 L/ha.

Grapevines: Where magnesium levels are low to marginal apply 3 L/ha at flower truss visible, flower buds separated, and at fruit set.

For grape stalk necrosis (grape shrivel) apply 3 applications of 4 to 5 L/ha commencing at pea-sized berries and again at veraison and 1 month before harvest. Note: For table grapes, applications should be completed 1 month before harvest. Water rate: 500-1000 L/ha.

Onion: 4 L/ha when there is sufficient leaf area to intercept spray. Repeat application as necessary. Water rate: 50-200 L/ha.

Potatoes: Where low to marginal levels exist, apply 2 applications of 4 L/ha between 1 to 4 weeks after 100% emergence. For improved dry matter, also apply during bulking. Water rate: 50-200 L/ha.

Poppy: One to three applications of 4 l/ha at 10 to 14 day intervals commencing once there is adequate leaf cover to intercept the spray and continuing up to the start of flowering. Water rate: 200 L/ha.

Tomato (Field Grown): 5 L/ha commencing when plants are 15 cm high and repeated 10 to 14 days later. Water rate: 50-500 L/ha.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS.

Mix the product thoroughly before use. Pour through the top filter into the tank three-quarters filled with water, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use and advice regarding whether or not product(s) should be co-applied. Many variables, outside the control of Yara, can influence the performance of co-applied products and therefore co-application is entirely at the risk of the end-user. It is strongly recommended that a limited application is made initially when using unfamiliar product mixes. *Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.*

PRECAUTIONS


Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. *Protect from frost and other climatic extremes. (temperatures exceeding 40°C)*

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY

Yara warrants that the product within the unopened container conforms to the chemical description on this label. The directions on this label must be followed carefully and are based on tests, believed to be reliable. As the occurrence of unusual weather or soil conditions, and the use of this product, are all beyond the control of Yara, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of Yara exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Goods are sold subject to the standard Conditions of Sale of Yara, available on request or at www.yaravita.com. Use of the product implies that the purchaser accepts these Conditions of Sale.



5