

# Mancozin

## RECOMMENDED APPLICATION RATES AND TIMINGS

**Acacia:** 2 to 3 l/ha applied during spring or autumn flush.

Water rate: 20 l/ha minimum.

**Beans, Peas:** 1 L/ha at the 4 to 6 leaf stage. Water rate: 200 L/ha

**Cereals:** 0.5-1 L/ha from 2 leaf stage to first node detectable (Zadoks G.S. 12 to 31). Water rate: 200 L/ha. For severe deficiency increase to 2 L/ha or repeat the above dose 10 to 14 days later.

**Citrus:** 1-2 L/ha during spring and autumn flushes.

Water rate: 500-2000 L/ha.

**Eucalyptus:** 2 to 3 l/ha applied during spring or autumn flush.

Water rate: 20 l/ha minimum.

**Grapevines:** 1.5 L/ha applied at 10 to 14 day intervals pre- and post-flowering. Avoid applications during flowering.

Water rate: 200-1000 L/ha.

**Lucerne:** 3.5 L/ha at dormancy break. Repeat after every cut when re-growth has occurred, as necessary. Water rate: 30-200 L/ha.

**Lupin:** 1 l/ha at the 4 to 6 leaf stage. Repeat once or twice at 10 to 14 day intervals if necessary. (For winter varieties apply the repeat sprays at spring re-growth). Water rate: 30 to 200 L/ha.

**Maize:** 1 l/ha at the 4 to 8 leaf stage. For moderate to severe deficiency repeat applications at 10 to 14 day intervals.

Water rate: 30-200 L/ha.

**Onion:** 3.5 L/ha when there is sufficient leaf area to obtain coverage (4 to 6 leaf). Repeat during bulking if necessary.

Water rate: 30-200 L/ha.

**Potatoes:** 1 L/ha at one week after 100% emergence and following petiole analysis during tuber bulking. For moderate to severe deficiency repeat applications at 10 to 14 day intervals.

Water rate: 30-250 L/ha.

## DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS.

*Mix the product thoroughly before use.* Pour through the top filter into the tank three-quarters filled with water, whilst agitating. Top up with water and continue to agitate until spraying is completed.

## TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use and advice regarding whether or not product(s) should be co-applied. Many variables, outside the control of Yara, can influence the performance of co-applied products and therefore co-application is entirely at the risk of the end-user. It is strongly recommended that a limited application is made initially when using unfamiliar product mixes. *Before co-application of products you, or your advisor, must visit [www.tankmix.com](http://www.tankmix.com) for important information. Alternatively, contact Yara for specific advice.*

## PRECAUTIONS


Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

## STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. *Protect from frost and other climatic extremes. (temperatures exceeding 40°C)*

## CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY

Yara warrants that the product within the unopened container conforms to the chemical description on this label. The directions on this label must be followed carefully and are based on tests, believed to be reliable. As the occurrence of unusual weather or soil conditions, and the use of this product, are all beyond the control of Yara, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of Yara exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Goods are sold subject to the standard Conditions of Sale of Yara, available on request or at [www.yaravita.com](http://www.yaravita.com). Use of the product implies that the purchaser accepts these Conditions of Sale.



5