

Molytrac Twin

WARNING

Excessive molybdenum can be harmful to stock. Do not apply fertilizer containing molybdenum on more than 25% of farm annually. Do not graze for four weeks after application.

RECOMMENDED APPLICATION RATES AND TIMINGS

Bananas: 0.25 L/ha. Regular applications may be necessary where low or marginal levels of molybdenum exist.

Water rate: 30 to 300 L/ha.

Beans, Peas: 250 to 500 mL/ha when plants are 5 to 15 cm tall.

For moderate to severe deficiency, repeat applications should be made at the above rate at 10 to 14 day intervals from the first application. Water rate: 30 to 200 L/ha.

Seed Dressing: 3 litres per tonne.

Brassicas: 300 to 500 mL/ha soil applied prior to emergence and at the 4 to 9 leaf stage. Water rate: 30 to 500 L/ha.

Canola: 250 to 500 mL/ha at 4 to 6 leaf stage and at onset of stem extension. Water rate: 30 to 500 L/ha.

Seed Dressing: 4 litres per tonne.

Capsicums (Field Grown): 0.25 L/ha 30 days after transplanting. Water rate: 70 L/ha.

Cereals: 125 to 250 mL/ha applied pre-planting to the soil with a herbicide where tank mixable. Water rate: 30 to 100 L/ha. Seed Dressing: 0.5 to 2 litres per tonne.

Citrus: 0.25 L/ha from 2/3 of new leaf development in the spring. For moderate to severe deficiency repeat application 10 to 14 days later. Also, 0.25 L/ha applied during autumn flush. Repeat 10 to 14 days later if necessary. Water rate: 500 L/ha.

Cotton: 250 to 500 mL/ha at the 4 to 6 leaf stage. Water rate: 30 to 200 L/ha.

Cucurbits (Field Grown): 250 mL/ha one to two weeks after germination and at the 4 leaf stage. Water rate: 200 L/ha

Grapevines: 0.5 L/ha applied before flowering. Repeat at fruit set, bunch closure and early veraison if necessary. Water rate: 200 L/ha

Lettuce (Field Grown): 250 to 500 mL/ha 10 to 14 days after transplanting or emergence. Repeat at 7 to 10 day intervals if necessary. Water rate: 50 to 500 L/ha.

Lucerne: 250 to 500 mL/ha every 2 to 3 cuts.

Water rate: 50 to 200 L/ha. Seed Dressing: 2 to 3 litres per tonne

Lupin: 250 to 500 mL/ha when the plants are 5 to 15 cm tall. Water rate: 50 to 200 L/ha.

Peanuts: 0.25 L/ha at the 4 to 6 leaf stage. Consider a second application 10 to 14 days later. Water rate: 30 to 200 L/ha.

Potatoes: 0.5 L/ha between 100% emergence and tuber initiation. Water rate: 50 to 200 L/ha.

Tomato (Field Grown): 250 mL/ha when crop is 15 cm tall. Water rate: 50 to 500 L/ha.

SOIL APPLICATION:

250 to 500 mL/ha soil applied pre-drilling, pre-planting or prior to emergence (or before regrowth of perennial crops).

Water rate: 30 to 200 L/ha.

For further crop recommendations visit Product Information at www.yara.com.au

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS.

Foliar/Soil application: *Mix the product thoroughly before use.* Pour through the top filter into the tank three-quarters filled with water, whilst agitating. Top up with water and continue to agitate until spraying is completed.

Seed Treatment: *Mix the product thoroughly before use.* Care should be taken to calibrate the machine accurately, to obtain an evenly applied correct dosage. Product application to most cereal and legume seed is enhanced by diluting them with water; this can vary with the moisture of the grain and the equipment used. Typically a total combined volume of seed treatment and water of 10 litres per tonne of seed is necessary for adequate coverage. Treatment of canola seed requires mixing equipment and the addition of water is not recommended.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use and advice regarding whether or not product(s) should be co-applied. Many variables, outside the control of Yara, can influence the performance of co-applied products and therefore co-application is entirely at the risk of the end-user. It is strongly recommended that a limited application is made initially when using unfamiliar product mixes. *Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.*

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE OF TREATED SEED


Treated seed should be drilled in season of application. Although the product should not have any effect on storage life of treated seed, the subsequent germination and performance of such seed may be adversely affected by physical storage conditions. No liability can therefore be accepted for the performance of stored, treated seed.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. *Protect from frost and other climatic extremes. (temperatures exceeding 40°C)*

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY

Yara warrants that the product within the unopened container conforms to the chemical description on this label. The directions on this label must be followed carefully and are based on tests, believed to be reliable. As the occurrence of unusual weather or soil conditions, and the use of this product, are all beyond the control of Yara, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of Yara exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Goods are sold subject to the standard Conditions of Sale of Yara, available on request or at www.yaravita.com. Use of the product implies that the purchaser accepts these Conditions of Sale.



5 033704 000545