

Phozin Twin with Synergise Technology

RECOMMENDED APPLICATION RATES AND TIMINGS

Acacia: 2 to 3 L/ha applied during spring or autumn flush.

Water rate: 20 L/ha minimum.

Beans: 1 to 2 L/ha when crop is 5 to 15 cm tall.

Water rate: 200 L/ha. Seed Dressing: 8 litres per tonne.

Canola: 1 to 2 L/ha at the 4 to 9 leaf stage.

Water rate: 200 to 500 L/ha. Seed Dressing: 15 litres per tonne.

Cereals: 1 to 2 L/ha from 2 leaf stage to first node detectable (Zadok's G.S 12 to 31), repeated once or twice at 10 to 14 day intervals in cases of moderate to severe deficiency.

Water rate: 70 L/ha. Seed Dressing: 6 litres per tonne.

Chick Peas: Seed Dressing: 8 litres per tonne.

Cotton: 1 to 2 L/ha 3 to 4 weeks after emergence. Repeat as required with 10 to 14 days between treatments.

Water rate: 70 L/ha.

Seed Dressing: Typical rate: Up to 18 litres per tonne if necessary.

Eucalyptus: 2 to 3 L/ha applied during spring or autumn flush.

Water rate: 20 L/ha minimum.

Maize: 1 to 2 L/ha from 3 to 8 leaf stage. For severe deficiency, repeat applications should be made at the above rate at 10 to 14 day intervals. Water rate: 200 L/ha. Seed Dressing: 8 litres per tonne.

Peas: 1 to 2 L/ha when crop is 5 to 15 cm tall.

Water rate: 200 L/ha. Seed Dressing: 8 litres per tonne.

Sorghum: 1 to 2 L/ha from 3 to 8 leaf stage. For severe deficiency repeat applications should be made at the above rate at 10 to 14 day intervals. Water rate: 70 L/ha.

Seed Dressing: Up to 18 litres per tonne if necessary.

SOIL APPLICATION

All Crops: 10 to 20 L/ha. Apply pre-drilling or pre-planting (or before regrowth of perennial crops). Water rate: 50 L/ha minimum.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS.

Foliar/Soil application: *Mix the product thoroughly before use.* Pour through the top filter into the tank three-quarters filled with water, whilst agitating. Top up with water and continue to agitate until spraying is completed.

Seed Treatment: *Mix the product thoroughly before use.* Care should be taken to calibrate the machine accurately, to obtain an evenly applied correct dosage. Product application to most cereal and legume seed is enhanced by diluting them with water; this can vary with the moisture of the grain and the equipment used. Typically a total combined volume of seed treatment and water of 10 litres per tonne of seed is necessary for adequate coverage. Treatment of canola seed requires mixing equipment and the addition of water is not recommended.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use and advice regarding whether or not product(s) should be co-applied. Many variables, outside the control of Yara, can influence the performance of co-applied products and therefore co-application is entirely at the risk of the end-user. It is strongly recommended that a limited application is made initially when using unfamiliar product mixes. *Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.*

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE OF TREATED SEED


Treated seed should be drilled in season of application. Although the product should not have any effect on storage life of treated seed, the subsequent germination and performance of such seed may be adversely affected by physical storage conditions. No liability can therefore be accepted for the performance of stored, treated seed.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. *Protect from frost and other climatic extremes. (temperatures exceeding 40°C)*

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY

Yara warrants that the product within the unopened container conforms to the chemical description on this label. The directions on this label must be followed carefully and are based on tests, believed to be reliable. As the occurrence of unusual weather or soil conditions, and the use of this product, are all beyond the control of Yara, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of Yara exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Goods are sold subject to the standard Conditions of Sale of Yara, available on request or at www.yaravita.com. Use of the product implies that the purchaser accepts these Conditions of Sale.



5 033704 002358

For further crop recommendations visit Product Information at www.yara.com.au