

# Seniphos

## RECOMMENDED APPLICATION RATES AND TIMINGS

**Apple, Pears:** 3 to 5 applications of 10-12.5 L/ha at 7 to 10 day intervals commencing 1 week after petal fall. Water rate: 500-1000 L/ha. Top Fruit Colour Enhancement: 10 L/ha in 500-1,000 L/ha water has been shown to enhance red colour development in apples, peaches and plums. Seniphos used for this purpose should not mixed with other agro-chemicals.

**Brassicas:** 2 to 3 applications of 10 L/ha from stem extension/head development at 7 to 10 day intervals. Water rate: 200-500 L/ha.

**Capsicums (Field Grown):** Up to 4 applications of 5 L/ha from flowering on second truss. Repeat at 10 to 14 day intervals. Water rate: 500 L/ha.

**Carrot:** 5 L/ha. 2 to 3 applications when crop is 15 cm tall and at 7 to 10 day intervals. Water rate: 200 L/ha.

**Citrus:** 10 L/ha at fruit set with 1 to 2 further applications at 7 to 10 day intervals. Water rate: 500-1000 L/ha.

**Cucumber, Cucurbits, Melon, Pumpkin, Squash, Water Melon, Zucchini (Field Grown):** 3 to 5 applications of 5 L/ha at 7 day intervals, commencing at first flowers through fruit expansion. Water rate: 200-500 L/ha.

**Grapevines:** 3 to 5 applications of 10 L/ha beginning at fruit set and repeated at 7 to 10 day intervals. Water rate: 500-1000 L/ha.

**Olive:** 10 L/ha before flowering. Repeat at 10 L/ha 10 to 14 days after flowering. Water rate: 200-1000 L/ha.

**Onion:** 1 to 2 applications of 5 L/ha during bulb filling. Water rate: 200-500 L/ha.

**Peach, Plum:** 3 applications of 10 to 12.5 L/ha commencing 3 to 7 days after petal fall and repeated at 7 to 10 day intervals. Water rate: 500-1000 L/ha. Top Fruit Colour Enhancement: 10 L/ha in 500-1000 L/ha water has been shown to enhance red colour development in apples, peaches and plums. Seniphos used for this purpose should not mixed with other agro-chemicals.

**Potatoes:** To increase tuber numbers, 1 application of 10 to 15 L/ha at tuber initiation (stolon hooking). To promote tuber size, 2 to 3 applications of 10 L/ha during bulking. Water rate: 200-500 L/ha.

**Tomato (Field Grown):** 2 to 4 applications of 5 L/ha commencing at flowering on second truss onwards. Repeat at 10 to 14 day intervals. Water rate: 200-500 L/ha.

## DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS.

*Mix the product thoroughly before use.* Pour through the top filter into the tank three-quarters filled with water, whilst agitating. Top up with water and continue to agitate until spraying is completed.

## TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use and advice regarding whether or not product(s) should be co-applied. Many variables, outside the control of Yara, can influence the performance of co-applied products and therefore co-application is entirely at the risk of the end-user. It is strongly recommended that a limited application is made initially when using unfamiliar product mixes. *Before co-application of products you, or your advisor, must visit [www.tankmix.com](http://www.tankmix.com) for important information. Alternatively, contact Yara for specific advice.*

## PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

## STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. *Protect from frost and other climatic extremes. (temperatures exceeding 40°C)*

## CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY

Yara warrants that the product within the unopened container conforms to the chemical description on this label. The directions on this label must be followed carefully and are based on tests, believed to be reliable. As the occurrence of unusual weather or soil conditions, and the use of this product, are all beyond the control of Yara, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of Yara exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Goods are sold subject to the standard Conditions of Sale of Yara, available on request or at [www.yaravita.com](http://www.yaravita.com). Use of the product implies that the purchaser accepts these Conditions of Sale.

