



YaraVita™ Agripotash

Potassium Fertiliser Solution

Guaranteed Analysis:

potassium (K)	41.5% w/v
---------------	-----------

Benefits:

- Formulated for safe application at critical growth stages to satisfy crop requirements
- Widely tank mixable with other crop sprays. Visit www.tankmix.com/yara for details
- Proven, reliable performance. Trialled and tested on a wide range of crops around the world
- High quality, consistent product. Manufactured to ISO 9001 quality assurance standards
- Easy to use liquid formulation. Pours and disperses easily and quickly into the spray tank
- High nutrient content means lower application rates reducing handling time and waste packaging

Key Nutrients:

- Potassium - Plays a key role in the transport of materials within the plant particularly to storage tissues like fruit, seeds and tubers. K also has a role in cell membrane structure, carbohydrate metabolism and energy accumulation and utilisation.



For details of product recommendations please see overleaf



Product Recommendations

Typical Crop Recommendations*

- **Beans:** 5 L/ha before flowering.
Water rate: 30-200 L/ha.
- **Beetroot:** 3 to 5 L/ha during bulb swelling.
Repeat applications at 10 to 14 day intervals if necessary. Water rate: 200 L/ha.
- **Brassicac:** 3-5 L/ha. Commence application at 4 to 6 true leaves and repeat as necessary at 7 to 10 day intervals between applications. Water rate: 250-750 L/ha.
- **Broccoli:** 3-5 L/ha. Commence application at 4 to 6 true leaves and repeat as necessary at 7 to 10 day intervals between applications. Water rate: 250-750 L/ha.
- **Brussel Sprouts:** 3-5 L/ha. Commence application at 4 to 6 true leaves and repeat as necessary at 7 to 10 day intervals between applications. Water rate: 250-750 L/ha.
- **Cabbage:** 3-5 L/ha. Commence application at 4 to 6 true leaves and repeat as necessary at 7 to 10 day intervals between applications. Water rate: 250-750 L/ha.
- **Calabrese:** 3-5 L/ha. Commence application at 4 to 6 true leaves and repeat as necessary at 7 to 10 day intervals between applications. Water rate: 250-750 L/ha.
- **Canola:** 5 L/ha at the 4 to 6 leaf stage.
Repeat as required for moderate to severe deficiency at 7 to 14 day intervals.
Water rate: 200 L/ha.
- **Carrot:** 3-5 L/ha. When sufficient leaf area to intercept spray with 10 to 14 day intervals between applications.
Water rate: 200-500 L/ha.
- **Cauliflower:** 3-5 L/ha. Commence application at 4 to 6 true leaves and repeat as necessary at 7 to 10 day intervals between applications. Water rate: 250-750 L/ha.
- **Cereals:** 5 L/ha from mid-tillering to 2nd node detectable (Zadoks G.S. 25 to 32). Repeat at 10 to 14 day intervals between these growth stages as necessary. Water rate: 200 L/ha. For aerial application: 3 L/ha from mid-tillering to 2nd node detectable (Zadoks G.S. 25 to 32). Repeat at 10 to 14 day intervals between these growth stages as necessary. Water rate: 20-50 L/ha.
- **Chick Peas:** 5 L/ha before flowering.
Water rate: 30-200 L/ha.
- **Citrus:** 3-5 L/ha. Commence applications after fruit set and repeat as necessary at 14 to 21 day intervals.
Water rate: At least 1000 L/ha.
- **Colton:** 2-5 L/ha. 3 applications at 14 day intervals commencing prior to first flower.
Water rate: 30-100 L/ha.
- **Eucalyptus:** 3 L/ha applied during spring or autumn flush. Water rate: 20 L/ha minimum.
- **Fertigation:** Up to 50 L/ha. Repeat the application at 7 to 14 day intervals as necessary.
- **Peanuts:** 5 L/ha before flowering.
Water rate: 200 L/ha.
- **Lentil:** 5 L/ha at the 4-6 leaf stage. For moderate to severe deficiency repeat the application 10-14 days later.
Water volume: 30-200 L/ha.
- **Lettuce (Field Grown):** 3 to 5 L/ha at 4 to 6 leaf stage with repeat applications at 7 to 10 day intervals as necessary.
Water rate: 500 L/ha.
- **Lucerne:** 5 L/ha. When sufficient leaf area to intercept spray. Water rate: 50 to 250 L/ha.
- **Maize:** 5 L/ha at the 4 to 8 leaf stage.
Water rate: 30-200 L/ha.
- **Onion:** 5 L/ha. When sufficient leaf area to intercept spray. Water rate: 200-500 L/ha.
- **Pineapple:** 5 L/ha. Apply 4 to 6 weeks before harvest. Water rate: 400-600 L/ha.
- **Potatoes:** 3 to 5 L/ha. Apply during bulking and repeat as necessary at 7 to 10 day intervals.
Water rate: 200-500 L/ha.
- **Tomato (Field Grown):** 3 to 5 L/ha. Commence applications at flowering of second truss and repeat as necessary at 7 to 10 day intervals.
Water rate: 200-500 L/ha.
- **Turf:** 10 L/ha. As soon as growth commences in the spring. In the case of moderate to severe deficiency, repeat applications at 14 day intervals.
Water rate: 500-1000 L/ha.

*The information provided is accurate to the best of Yara's knowledge and belief. Any recommendations are meant as a guide and must be adapted to suit local conditions. Always read the label before use.



YARA AUSTRALIA PTY LTD
201 Miller Street, North Sydney, NSW 2060
Australia
Freecall: 1800 684 266
Tel.: 02 9959 4266 Fax 02 9959 4050
Websites: www.yara.com.au www.yaraphosyn.com

Phosyn Analytical (& MEGALAB)
Tel.: 07 5568 8700 Fax 07 5522 0720
Email phosynanalytical@phosyn.com

