

Stopit

RECOMMENDED APPLICATION RATES AND TIMINGS

Apple, Pears: Repeat applications of 7.5-10 L/ha applied from end of flowering to harvest. Refer to YARA fruit quality programme for precise timings. Water rate: 500-1000 L/ha. When using in a tank mix with Seniphos the higher water rate must be used.

Apricot, Peach: 2 applications of 5 L/ha followed by 2 applications of 10 L/ha starting 3 to 7 days after petal fall and repeated at 7 to 10 day intervals. Plus 10 L/ha applied 14 to 21 and again 7 to 10 days before harvest. Water rate: 500-1000 L/ha.

Bananas: Four to five applications of 5 L/ha from fruit set at 10 to 14 day intervals. Water rate: 500 L/ha.

Brassicas: 2 to 3 applications of 5 L/ha applied from stem extension/head development to harvest with 10 to 14 day intervals between applications. Water rate: 200-500 L/ha.

Capsicum, Pepper, Tomato (Field Grown): 1 to 4 applications of 5 L/ha commencing at flowering of second truss. Allow 7 days between applications. Water rate: 50-500 L/ha.

Cherry, Plum: 10 L/ha applied 1 month and again 2 weeks before harvest. Water rate: 500-1000 L/ha.

Citrus: 3-5 applications of 20 L/ha applied at 10 to 14 day intervals commencing when fruit reach 40 to 50 mm diameter (around mid January for Washington Navels in the Riverland and Sunraysia). Water rate: 1000 L/ha minimum.

Cucumber, Cucurbits, Melon, Pumpkin, Squash, Water Melon, Zucchini (Field Grown): 2 to 3 applications of 5 L/ha commencing at fruit set with 7 to 10 day intervals between applications. Water rate: 200-500 L/ha.

Grapevines: 5 to 10 L/ha at 7 to 10 day intervals from fruit set to one month before harvest. Water rate: 500-1000 L/ha. Water rate: 500-1000 L/ha.

Lettuce (Field Grown): 2 to 3 applications of 5 L/ha commencing 10 to 14 days after transplanting or emergence with 7 to 10 day intervals between applications. Water rate: 50-500 L/ha.

Mango: 3 to 5 applications of 5 L/ha applied during the period from early fruiting to fruit swelling. Allow 7 to 10 days between applications. Water rate: 500-1000 L/ha.

Nectarines, Olives: Regular applications of 5 to 10 L/ha at 7 to 14 day intervals from petal fall to one week before harvest. Water rate: 150-1000 L/ha.

Onions: 5 to 10 L/ha at bulb swelling. Repeat applications at 7 to 10 day intervals as necessary. Water rate: 200-500 L/ha.

Poppy: One to three applications of 5 L/ha at 10 to 14 day intervals commencing once there is adequate leaf cover to intercept the spray and continuing up to the start of flowering. Water rate: 200 L/ha.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS.

Mix the product thoroughly before use. Pour through the top filter into the tank three-quarters filled with water, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use and advice regarding whether or not product(s) should be co-applied. Many variables, outside the control of Yara, can influence the performance of co-applied products and therefore co-application is entirely at the risk of the end-user. It is strongly recommended that a limited application is made initially when using unfamiliar product mixes. *Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.*

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. *Protect from frost and other climatic extremes. (temperatures exceeding 40°C)*

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY

Yara warrants that the product within the unopened container conforms to the chemical description on this label. The directions on this label must be followed carefully and are based on tests, believed to be reliable. As the occurrence of unusual weather or soil conditions, and the use of this product, are all beyond the control of Yara, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of Yara exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Goods are sold subject to the standard Conditions of Sale of Yara, available on request or at www.yaravita.com. Use of the product implies that the purchaser accepts these Conditions of Sale.

