

Zinphos

RECOMMENDED APPLICATION RATES AND TIMINGS

Beans, Soya Bean: 3 L/ha when crop is 5 to 15 cm tall, repeated at 10 to 14 day intervals if necessary.

Water rate: 30-100 L/ha maximum.

Cereals: 1-2 L/ha. Apply from 4 leaf stage to early tillering.

Repeat 10 to 14 days later if necessary. Water rate: 30-100 L/ha.

Chick Peas: 2.5 L/ha at 10 to 15 cm stage.

Water rate: 30-100 L/ha maximum.

Cotton: 2-5 L/ha. Apply at 4 to 6 leaf stage and at green bolls over 50 mm long. Water rate: 50-100 L/ha.

Peanuts: 3 L/ha at the 4 to 6 leaf stage. Water rate: 30-100 L/ha.

Lucerne: 3 L/ha applied to early season regrowth. Consider a second application 2 to 3 weeks later.

Water rate: 30-100 L/ha maximum.

Maize: 2 to 4 L/ha. Apply at 4 to 8 leaf stage.

Water rate: 30-100 L/ha.

Peas: 2.5 L/ha at 10 to 15 cm stage. For moderate to severe deficiency, a repeat application may be necessary 10 to 14 days later. Water rate: 30-100 L/ha maximum.

Potatoes: Apply 3 to 5 L/ha at the following timings:

1st: One week after 100% emergence.

2nd: To enhance maximum viable tuber number, apply at tuber initiation.

3rd: To help to increase tuber size, repeat applications at 10 to 14 day intervals as soon as tubers are 10 mm in diameter (minimum 2 applications). Water rate: 30-100 L/ha maximum.

For aerial applications 2 L/ha applied at the timings described above. Water rate: 20-50 L/ha.

Rice: 3 L/ha. Apply at start of tillering (when plants have 4 to 6 leaves). A repeat application can be made 10 days later.

Water rate: 50-100 L/ha.

Sunflower: 3 L/ha at the 3 to 4 leaf stage.

Water rate: 50-100 L/ha maximum.

SOIL APPLICATION

Maize: 20 L/ha applied at sowing using suitable equipment to direct neat product over the seed along the drill line. Water rate: nil

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS.

Mix the product thoroughly before use. Pour through the top filter into the tank three-quarters filled with water, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use and advice regarding whether or not product(s) should be co-applied. Many variables, outside the control of Yara, can influence the performance of co-applied products and therefore co-application is entirely at the risk of the end-user. It is strongly recommended that a limited application is made initially when using unfamiliar product mixes. *Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.*

PRECAUTIONS


Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. *Protect from frost and other climatic extremes. (temperatures exceeding 40°C)*

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY

Yara warrants that the product within the unopened container conforms to the chemical description on this label. The directions on this label must be followed carefully and are based on tests, believed to be reliable. As the occurrence of unusual weather or soil conditions, and the use of this product, are all beyond the control of Yara, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of Yara exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Goods are sold subject to the standard Conditions of Sale of Yara, available on request or at www.yaravita.com. Use of the product implies that the purchaser accepts these Conditions of Sale.



5